Pemetrexed (Alimta®)

Pronounced: pem-e-TREX-ed

Classification: Antifolate Antineoplastic Agent

About Pemetrexed

Pemetrexed works by blocking three separate enzymes that cells need to replicate. These enzymes, needed for folate-dependent metabolic processes, are required for DNA replication. Therefore, inhibiting these enzymes inhibits cellular growth.

How to Take Pemetrexed

Pemetrexed is given by intravenous (IV) infusion. It is typically given once every 21 days. The dose is based on the patient's height and weight. A steroid medication is often given before pemetrexed to decrease skin rash. Folic acid supplements should be given for at least 5 out of 7 days prior to the first dose of pemetrexed, throughout treatment and for three weeks after the last dose is given. An injection of vitamin B12 should be given before the first dose of pemetrexed, and then about every 9 weeks during treatment. These vitamin supplementations will decrease the side effects you experience. Speak with your provider concerning the specific dosing of steroids, folic acid and B12 and timing of these medications.

Let your healthcare provider know all medications, vitamins and supplements you are taking as some can interfere with pemetrexed. Be sure to notify your care provider if you take any form of NSAID (non-steroidal anti-inflammatory drug) such as aspirin or ibuprofen because, taken in conjunction with pemetrexed, can cause kidney problems.

Possible Side Effects of Pemetrexed

There are a number of things you can do to manage the side effects of pemetrexed. Talk to your doctor or nurse about these recommendations. They can help you decide what will work best for you. These are some of the most common side effects:

Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your WBC count can drop, putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

Tips to preventing infection:

- Washing hands, both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent mouth care.
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your doctor or nurse before scheduling dental appointments or procedures.
- Ask your doctor or nurse before you, or someone you live with, has any vaccinations.

Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the red cell count is low, you may feel tired or weak. You should let your doctor or nurse know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion.

Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the count is low you are at a higher risk of bleeding. Let your doctor or nurse know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin®, Aleve®, Advil®, etc. as these can all increase the risk of bleeding. Unless your healthcare team tells you otherwise, you may take acetaminophen (Tylenol).
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

Nausea and/or Vomiting

Talk to your doctor or nurse so they can prescribe medications to help you manage nausea and vomiting. In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try antacids, (e.g. milk of magnesia, calcium tablets such as Tums), saltines, or ginger ale to lessen symptoms.

Call your doctor or nurse if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

Decrease in Appetite

Nutrition is an important part of your care. Cancer treatment can affect your appetite and, in some cases, the side effects of treatment can make eating difficult. Ask your nurse about nutritional counseling services at your treatment center to help with food choices.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.
- You may experience a metallic taste or find that food has no taste at all. You may dislike foods or beverages that you liked before receiving cancer treatment. These symptoms can last for several months or longer after treatment ends.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products and fish without a strong smell. Sometimes cold food has less of an odor.
- Add extra flavor to meat or fish by marinating it in sweet juices, sweet and sour sauce or dressings. Use seasonings like basil, oregano or rosemary to add flavor. Bacon, ham and onion can add flavor to vegetables.

Fatigue

Fatigue is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

Diarrhea

Your oncology team can recommend medications to relieve diarrhea. Also, try eating low-fiber, bland foods, such as white rice and boiled or baked chicken. Avoid raw fruits, vegetables, whole grain breads, cereals and seeds. Soluble fiber is found in some foods and absorbs fluid, which can help relieve diarrhea. Foods high in soluble fiber include: applesauce, bananas (ripe), canned fruit, orange sections, boiled potatoes, white rice, products made with white flour, oatmeal, cream of rice, cream of wheat, and farina. Drink 8-10 glasses on non-alcoholic, un-caffeinated fluid a day to prevent dehydration.

Mouth Ulcers (Sores)

Certain cancer treatments can cause sores or soreness in your mouth and/or throat. Notify your doctor or nurse if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated or painful. Performing regular mouth care can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon salt in an eight ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages and citrus juices.

Skin Changes/Rash

These include dryness, itching, and rash. You should use a moisturizer on your skin and lips, but avoid moisturizers with perfumes or scents. Your doctor or nurse can recommend

medication if itching is bothersome. If your skin does crack or bleed, be sure to keep the area clean to avoid infection. Rash related to this medication typically occurs between doses and resolves before the next dose. A rash can result in blistering or peeling of the skin. Notify your oncology team if you develop a rash.

Other Side Effects

Some less common side effects that have been reported include: shortness of breath, blood clots, and an allergic reaction to the medication. Report any side effects to your healthcare provider.

Reproductive Concerns

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment. Even if your menstrual cycle stops or you believe you are not producing sperm, you could still be fertile and conceive. A woman should stop breastfeeding while on this medication because it is unknown if the medication can be passed through breast milk.

If you have questions or concerns about the medication that you have been prescribed, please contact your healthcare team. OncoLink is designed for educational purposes only and is not engaged in rendering medical advice or professional services. The information provided through OncoLink should not be used for diagnosing or treating a health problem or a disease. It is not a substitute for professional care. If you have or suspect you may have a health problem, you should consult your health care provider.

Carboplatin (Paraplatin®)

Pronounced: car-boe-PLATT-in
Classification: Platinum Chemotherapies

About Carboplatin

Carboplatin is a heavy metal compound that inhibits synthesis of RNA, DNA, and protein in cells. All of these compounds are vital for cells to divide and grow. By preventing them from dividing, the medication can stop the cancer from growing.

How to Take Carboplatin

Carboplatin is given by intravenous (IV, into a vein) injection. The schedule and dosage is based on the person's size and the cancer type being treated. It can be given alone, or with other drugs.

Carboplatin can interact with certain medications including some antibiotics, diuretics and blood thinners. Be sure to tell your healthcare provider about all medications and supplements you take.

You or anyone you live with should avoid having live (oral polio, measles, rotovirus, yellow fever) or live-attenuated vaccines while receiving this medication.

Possible Side Effects

There are a number of things you can do to manage the side effects of carboplatin. Talk to your doctor or nurse about these recommendations. They can help you decide what will work best for you. These are some of the most common side effects:

Nausea and/or Vomiting

Talk to your doctor or nurse so they can prescribe medications to help you manage nausea and vomiting. In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try antacids, (e.g. milk of magnesia, calcium tablets such as Tums), saltines, or ginger ale to lessen symptoms.

Call your doctor or nurse if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your WBC count can drop, putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

Tips to preventing infection:

- Washing hands, both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent mouth care.
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your doctor or nurse before scheduling dental appointments or procedures.
- Ask your doctor or nurse before you, or someone you live with, has any vaccinations.

Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the red cell count is low, you may feel tired or weak. You should let your doctor or nurse know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion.

Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the count is low you are at a higher risk of bleeding. Let your doctor or nurse know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin®, Aleve®, Advil®, etc. as these can all increase the risk of bleeding. Unless your healthcare team tells you otherwise, you may take acetaminophen (Tylenol).
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

Decrease in Appetite

Nutrition is an important part of your care. Cancer treatment can affect your appetite and, in some cases, the side effects of treatment can make eating difficult. Ask your nurse about nutritional counseling services at your treatment center to help with food choices.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.
- You may experience a metallic taste or find that food has no taste at all. You may dislike foods or beverages that you liked before receiving cancer treatment. These symptoms can last for several months or longer after treatment ends.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products and fish without a strong smell. Sometimes cold food has less of an odor.

Allergic Reactions

In some cases, patients can have an allergic reaction to this medication. Signs of a reaction can include: rash, itching, hives, flushing, and/or shortness of breath or difficulty breathing. If you notice any changes in how you feel during the infusion, let your nurse know immediately. The infusion will be slowed or stopped if this occurs. Depending on the severity of your reaction, you may still be able to receive the medication with a pre-medication to prevent a reaction, or if the medication is given at a slower rate.

Fatigue

Fatigue is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become thin, brittle, or may fall out. This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

Peripheral Neuropathy (Numbness or Tingling in the Hands and/or Feet)

Peripheral neuropathy is a toxicity that affects the nerves. It causes a numbness or tingling feeling in the hands and feet, often in the pattern of a stocking or glove. This can get progressively worse with additional doses of the medication. In some people, the symptoms slowly resolve after the medication is stopped, but for some it never goes away completely. You should let your healthcare provider know if you experience numbness or tingling in the hands and feet, as they may need to adjust the doses of your medication.

Mouth Sores

Certain cancer treatments can cause sores or soreness in your mouth and/or throat. Notify your doctor or nurse if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated or painful. Performing regular mouth care can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon salt in an eight ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages and citrus juices.

Diarrhea

Your oncology team can recommend medications to relieve diarrhea. Also, try eating low-fiber, bland foods, such as white rice and boiled or baked chicken. Avoid raw fruits, vegetables, whole grain breads, cereals and seeds. Soluble fiber is found in some foods and absorbs fluid, which can help relieve diarrhea. Foods high in soluble fiber include: applesauce, bananas (ripe), canned fruit, orange sections, boiled potatoes, white rice, products made with white flour, oatmeal, cream of rice, cream of wheat, and farina. Drink 8-10 glasses on non-alcoholic, un-caffeinated fluid a day to prevent dehydration.

Liver Toxicity

This medication can cause liver toxicity, which your doctor may monitor for using blood tests called liver function tests. If you develop elevations in your liver function tests, your healthcare provider may need to lower your dose or stop the medication. Notify your healthcare provider if you notice yellowing of the skin or eyes, your urine appears dark or brown or pain in your abdomen, as these can be signs of liver toxicity.

Nail Changes

Your fingernails/toenails may become dark, brittle or fall off. They may develop ridges or markings on them. Keep your fingernails and toenails clean and dry. You may use nail polish, but do not wear fake nails. If any nails fall off, clean the nail bed well with soap and water and cover with a bandaid.

Secondary Malignancies

There is a very low risk of developing another type of cancer due to treatment with this medication, which can occur many years after treatment.

Electrolyte Changes

Carboplatin can cause changes in the levels of certain electrolytes in your blood, including sodium, potassium, calcium and magnesium. Your healthcare team will monitor your electrolyte levels throughout treatment.

Kidney Problems

Carboplatin can impact your kidney function. Your healthcare team will monitor your kidney function throughout treatment. Intravenous fluids are given with the chemotherapy to flush your kidneys and increase your urine output. Try to drink at least 6-8 glasses of uncaffienated fluids a day. Call your doctor or nurse if you do not urinate for more than 12 hours.

Other Side Effects

In rare cases, this medication can cause temporary changes to hearing and vision. Contact your care team if you notice ringing in your ears, decrease in hearing, or changes in your vision.

Reproductive Concerns

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment. Even if your menstrual cycle stops or you believe you are not producing sperm, you could still be fertile and conceive. You should consult with your healthcare team before breastfeeding while receiving this medication.

If you have questions or concerns about the medication that you have been prescribed, please contact your healthcare team. OncoLink is designed for educational purposes only and is not engaged in rendering medical advice or professional services. The information provided through OncoLink should not be used for diagnosing or treating a health problem or a disease. It is not a substitute for professional care. If you have or suspect you may have a health problem, you should consult your health care provider.