

## Bleomycin (Blenoxane®)

**Pronounce:** blee-oh-MY-sin

**Classification:** Antineoplastic Antibiotic

### About Bleomycin (Blenoxane®)

The way bleomycin works is not fully understood. It is thought to affect cell reproduction and growth, reducing the number of cancer cells in the body.

### How to Take Bleomycin

Bleomycin is given by intravenous (IV, into a vein) infusion, intramuscular (IM, into a muscle), or subcutaneous (SQ, under the skin) injections. It can be given alone or in combination with other medications.

Bleomycin can also be used as a "sclerosing" agent to treat pleural effusions. In this case, it causes scarring of the pleural space to prevent the effusion (fluid) from building back up.

### Possible Side Effects

There are a number of things you can do to manage the side effects of bleomycin. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

#### Pulmonary (Lung) Toxicity

Bleomycin may cause serious lung problems. The risk may be higher in the elderly or those who received high doses, but lung effects have also happened in young patients and with low doses. There is a lifetime maximum dose of this medication due to the possibility of lung problems, which can include [pneumonitis](#) (inflammation of lung tissue) and pulmonary fibrosis (scarring and stiffening of the lung tissue). These problems can begin months to a few years after treatment is done. You may have breathing tests (pulmonary function tests, or PFTs) before starting bleomycin. Tell your healthcare provider if you have any shortness of breath, cough, or wheezing. If you are having a hard time breathing, call 911 or go to the closest emergency room right away.

If you are planning for any type of surgery, tell your healthcare team that you have received bleomycin, as there is a greater risk for having pulmonary (lung) problems from oxygen given during surgery.

It is strongly recommended that you do not smoke if you have received bleomycin, as this increases the risk of lung complications. You should report any shortness of breath, difficulty breathing, cough, or wheezing to your care provider. Lung complications can happen several years after treatment with bleomycin, so you should be sure all of your care providers know you received this medication and have any new lung symptoms checked. Of note, patients who have received this medication are recommended to have clearance by a dive medicine specialist prior to scuba diving.

#### Allergic-Like Reactions

An allergic-like reaction with fever, chills, low blood pressure, wheezing, or difficulty breathing may occur immediately or up to a few days after receiving treatment. If needed, medications are given to help with these effects. If you have any of these effects, tell your care team right away.

## Nail and Skin Changes

Your fingernails/toenails may become dark, thick, brittle, or fall off. Your skin may be dry, feel thick, or appear darker (hyperpigmentation). Your skin may be more sensitive to the sun, which can cause severe sunburn or rash. Sun sensitivity can last even after you have stopped taking the medication. Avoid the sun between 10 am to 2 pm when it is strongest. Wear sunscreen (at least SPF 15) every day, wear sunglasses and long sleeves/pants to protect your skin. Keep your fingernails and toenails clean and dry. You may use nail polish, but do not wear fake nails. Tell your provider if any nails fall off.

This medication can also cause radiation recall. This is redness, swelling, or blistering of the skin in an area that was previously treated (even years ago) with radiation. The goal of treatment for radiation recall is to manage the symptoms until it heals. Topical steroids or anti-inflammatory medications or cool compresses may help. Avoid sun exposure and tight-fitting clothes that would rub on the area.

## Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become [thin, brittle, or may fall out](#). This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats, and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

## Mouth Ulcers (Mucositis)

Certain cancer treatments can cause sores or soreness in your mouth and/or throat. Notify your oncology care team if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated, or painful. Performing [regular mouth care](#) can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt with warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon of salt in an eight-ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages, and citrus juices.

## Decrease in Appetite

[Nutrition](#) is an important part of your care. Cancer treatment can affect your appetite and, in some cases, the side effects of treatment can make eating difficult. Ask your nurse about nutritional counseling services at your treatment center to help with food choices.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.
- You may experience a metallic taste or find that food has no taste at all. You may dislike foods or beverages that you liked before receiving cancer treatment. These symptoms can last for several months or longer after treatment ends.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products, and fish without a strong smell. Sometimes cold food has less of an odor.
- Add extra flavor to meat or fish by marinating it in sweet juices, sweet and sour sauce, or dressings. Use seasonings like basil, oregano, or rosemary to add flavor. Bacon, ham, and onion can add flavor to vegetables.

## Nausea and/or Vomiting

Talk to your care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try antacids, (e.g. milk of magnesia, calcium tablets such as Tums), saltines, or ginger ale to lessen symptoms.

Call your provider if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

## Reproductive Concerns

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment. Even if your menstrual cycle stops or you believe you are not producing sperm, you could still be fertile and conceive. You should not breastfeed while receiving bleomycin.

# Cyclophosphamide IV Formulation (Cytoxan®, Neosar®, Endoxan®)

**Pronounce:** SYE-kloe-FOS-fa-mide

**Classification:** Alkylating Agent

## About Cyclophosphamide IV Formulation (Cytoxan®, Neosar®, Endoxan®)

Cyclophosphamide kills cancer cells through a process called alkylation. Alkylation damages the DNA of cells, which prevents them from dividing and causes them to die. Since cancer cells tend to divide faster and with less error correcting than healthy cells, cancer cells are more sensitive to this damage.

## How to Take Cyclophosphamide

Cyclophosphamide can be given by intravenous (IV, into a vein) infusion or taken orally (by mouth) in a pill form. This information is about the IV formulation. Your dose and schedule are determined by your size and type of cancer being treated.

This medication can interact with a number of medications including: metronidazole, tamoxifen, warfarin, cyclosporine amiodarone, Echinacea, and thiazide diuretics, among others. Be sure to tell your healthcare provider about all medications and supplements you take.

## Possible Side Effects

There are a number of things you can do to manage the side effects of cyclophosphamide. Talk to your care team about these recommendations. They can help you decide what will work best for you. There are some of the most common or important side effects:

### Infection and Low White Blood Cell Count (Leukopenia or Neutropenia)

This medication can cause life threatening infections, with or without a decrease in white blood cell counts.

White blood cells (WBC) are important for fighting infection. While receiving treatment, your [WBC count can drop](#), putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4°F or 38°C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

### Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bathe daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with, has any vaccinations.

## Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion.

## Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib) etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding use of these agents and all over the counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

## Nausea and/or Vomiting

Talk to your oncology care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

## Diarrhea

Your oncology care team can recommend medications to relieve [diarrhea](#). Also, try eating [low-fiber](#), bland foods, such as white rice and boiled or baked chicken. Avoid raw fruits, vegetables, whole grain breads, cereals and seeds. Soluble fiber is found in some foods and absorbs fluid, which can help relieve diarrhea. Foods high in soluble fiber include: applesauce, bananas (ripe), canned fruit, orange sections, boiled potatoes, white rice, products made with white flour, oatmeal, cream of rice, cream of wheat, and farina. Drink 8-10 glasses of non-alcoholic, un-caffeinated fluid a day to prevent dehydration.

## Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become [thin, brittle, or may fall out](#). This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The

use of scarves, wigs, hats and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

## Nail and Skin Changes

Your fingernails/toenails may become dark, brittle or fall off. You may notice dry skin or changes in the color or tone of your skin. Your skin may be more sensitive to the sun, which can result in severe sunburn or rash. Sun sensitivity can last even after chemotherapy is completed. Avoid the sun between 10-2pm, when it is strongest. Wear sunscreen (at least SPF 15) everyday, wear sunglasses and long sleeves/pants to protect your skin. Keep your fingernails and toenails clean and dry. You may use nail polish, but do not wear fake nails. If any nails fall off, clean the nail bed well with soap and water and cover with a band aid.

## Important but Less Common Side Effects

- **Bladder Irritation:** Cyclophosphamide may irritate your bladder and can cause blood in your urine or pain with urination. Drink at least 6 to 8 glasses of fluid per day to flush out your bladder. You may be given IV fluids along with this medication to help lower the chance of bladder irritation. Let your care team know if you have a hard time urinating, have pain or pink/red urine, or bleed with urination. When given in high doses, your provider may give you a medication to protect your bladder.
- **Secondary Malignancies:** There is a very low risk of developing leukemia or another type of cancer due to treatment with this medication, which can happen many years after treatment. This is most often linked to repeated treatments or high doses.
- **Electrolyte Abnormalities:** This medication can affect the normal levels of electrolytes (sodium, potassium, magnesium, calcium, etc.) in your body. Your levels will be monitored using blood tests. If your levels become too low, your care team may prescribe specific electrolytes to be given by IV or taken by mouth. Do not take any supplements without first consulting with your care team.
- **Lung Changes:** This medication may cause lung changes, including pneumonitis (irritation of the lung tissue) and pulmonary fibrosis (a scarring and stiffening of the lung tissue). These problems can develop during treatment, or months to years after treatment is completed. Call your provider right away if you have new or worsening shortness of breath, cough, wheezing or difficulty breathing.
- **Heart Problems:** This medication can affect your heart function and can cause abnormal heartbeats or an abnormal heart rhythm called QT prolongation. Call your provider right away if you develop swelling of the feet or ankles, shortness of breath, have rapid weight gain, feel abnormal heartbeats, or if you feel dizzy or faint.
- **Wound Healing:** This medication can lead to slower or incomplete wound healing, such as a surgical wound not healing or staying closed. Be sure to inform the team performing the surgical procedure that you are taking this drug. You should also inform your oncology team that a surgical procedure is planned. It is recommended that this medication be discontinued prior to any surgery. In addition, any surgical incision should be fully healed prior to starting or restarting the medication. If you have a surgical wound that has not healed or begins to have signs of infection (redness, swelling, warmth), report this to your healthcare team.
- **Veno-occlusive Liver Disease:** This medication can cause blood clots in the small veins of your liver. This can lead to liver failure. If you feel like there is fluid in your belly or notice that your skin or the whites of your eyes are yellowing, contact your care team right away.

## Reproductive Concerns

This medication may affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and [vaginal dryness](#). In addition, the desire for sex may decrease during treatment. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your

oncology team.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. For women, effective birth control is necessary during treatment and for at least 1 year after treatment. For men, effective birth control is necessary during treatment and for at least 4 months after treatment. Even if your menstrual cycle stops or you believe you are not producing sperm, you could still be fertile and conceive. You should not breastfeed during treatment and for at least 1 week after your last dose.

## Doxorubicin (Adriamycin®, Rubex®)

**Pronounce:** DOX-oh-ROO-bi-sin

**Classification:** Anthracycline

### About Doxorubicin (Adriamycin®, Rubex®)

Anthracyclines work by interrupting the copying of DNA, which is necessary for cancer cell growth. This causes the cancer cells to die, slowing or stopping tumor growth. Doxorubicin interferes with the growth of cancer cells and slows their spread in the body by inhibiting DNA synthesis and causing the production of harmful free radicals.

### How to Take Doxorubicin

Doxorubicin is given through an intravenous (IV, into a vein) infusion or injection. It can be given alone or with other drugs. The dosage and schedule are determined by your height and weight, type of cancer, and how the medication is given.

Even when carefully and correctly administered by trained personnel, this drug may cause a feeling of burning and pain. There is a risk that this medication may leak out of the vein at the injection site, leading to tissue damage that can be severe. If the area of injection becomes red, swollen, or painful at any time during or after the injection, tell your care team right away. Do not apply anything to the site unless told to do so by your care team.

This medication is red. Your urine may look orange or reddish in color for 1-2 days after the infusion. This is not blood. This is expected as the medication is cleared from your body. If the red urine lasts more than two days or if you have other urinary symptoms, such as frequency or painful urination, call your healthcare provider.

The blood levels and effectiveness of this medication can be affected by certain foods and medications, so they should be avoided. These include verapamil, phenytoin, fluconazole, voriconazole, St. John's wort, phenobarbital, trastuzumab, dexrazoxane, and 6-mercaptopurine, among others. Be sure to tell your healthcare provider about all medications and supplements you take.

### Possible Side Effects of Doxorubicin

There are a number of things you can do to manage the side effects of doxorubicin. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

#### Heart Problems

In rare cases, the heart muscle can be damaged by this medication, causing heart failure and cardiomyopathy. This heart damage can occur during therapy or many months to years after treatment. The risk is highest at higher doses, in patients who receive other cardio-toxic medications, radiation to the chest area, and in children. There is a maximum lifetime dose that you can receive of this medication. Your provider may order tests to check your heart function before you begin treatment or if any symptoms arise.

If you have shortness of breath, new or worsening cough, ankle swelling, chest pain, rapid or irregular heartbeats, call your provider right away, or call 911.

## Secondary Cancers

A secondary cancer is one that develops as a result of cancer treatment for another cancer. This is quite rare, but you should be aware of the risk. In most cases, a secondary cancer related to chemotherapy is a blood cancer (leukemia, lymphoma). This can occur years after treatment. This is most often associated with repeated treatments or high doses. Your provider will monitor your labs closely. You may need a complete blood count with differential checked each year by your healthcare provider if you received high-risk therapies.

## Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your [WBC count can drop](#), putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4°F or 38°C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

### Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever, or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bathe daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with has any vaccinations.

## Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing, or pain in your chest. If the count gets too low, you may receive a blood transfusion.

## Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums, or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib), etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding the use of these agents and all over-the-counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

## Loss or Thinning of Scalp and Body Hair (Alopecia)



Your hair may become [thin, brittle, or may fall out](#). This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats, and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

## Nausea and/or Vomiting

Talk to your oncology care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

## Diarrhea

Your oncology care team can recommend medications to relieve [diarrhea](#). Also, try eating [low-fiber](#), bland foods, such as white rice and boiled or baked chicken. Avoid raw fruits, vegetables, whole-grain breads, cereals, and seeds. Soluble fiber is found in some foods and absorbs fluid, which can help relieve diarrhea. Foods high in soluble fiber include applesauce, bananas (ripe), canned fruit, orange sections, boiled potatoes, white rice, products made with white flour, oatmeal, cream of rice, cream of wheat, and farina. Drink 8-10 glasses of non-alcoholic, un-caffeinated fluid a day to prevent dehydration.

## Mouth Ulcers (Mucositis)

Certain cancer treatments can cause [sores or soreness in your mouth](#) and/or throat. Notify your oncology care team if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated or painful. Performing [regular mouth care](#) can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt with warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon of salt in an eight-ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages, and citrus juices.

## Nail and Skin Changes

Your fingernails/toenails may become dark, brittle, or fall off. You may notice dry skin or changes in the color or tone of your skin. Your skin will be more sensitive to the sun, which can result in severe sunburn or rash. Sun sensitivity can last even after chemotherapy is completed. Avoid the sun between 10-2 pm, when it is strongest. Wear sunscreen (at least SPF 15) every day, wear sunglasses and long sleeves/pants to protect your skin. Keep your fingernails and toenails clean and dry. You may use nail polish, but do not wear fake nails. If any nails fall off, clean the nail bed well with soap and water and cover with a bandaid.

## Important but Less Common Side Effects

- **Radiation Recall:** Radiation recall is when the administration of a medication causes a skin reaction that looks like a sunburn (redness, swelling, soreness, peeling skin) in areas where radiation was previously given. Notify your oncology team if you notice this side effect. Treatment can include topical steroid ointments and a delay in your next chemotherapy dose.
- **Tumor Lysis Syndrome:** If there are a large amount of tumor cells in your body prior to treatment, you are at risk for tumor lysis syndrome. This happens when the tumor cells die too quickly and their waste



overwhelms the body. You may be given a medication (allopurinol) and IV fluids to help prevent this. If you experience nausea, vomiting, diarrhea or become lethargic (drowsy, sluggish), notify your oncology team right away. TLS can affect your kidney function. Your provider will monitor your kidney function with blood work. Notify your provider if you have little or no urine output.

## Sexual & Reproductive Concerns

This drug may affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and [vaginal dryness](#). In addition, the desire for sex may decrease during treatment.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Women should use effective birth control during and for 6 months after treatment, even if your menstrual cycle stops. Men should use condoms during and for three months after treatment even if you believe you are not producing sperm. If you have a pregnant partner, you should use condoms during and for 10 days after the last dose. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team. You should not breastfeed while receiving this medication.

## Etoposide (Toposar®, VePesid®, Etopophos®, VP-16)

**Pronounce:** e-toe-POE-side

**Classification:** Topoisomerase inhibitor

### About Etoposide (Toposar®, VePesid®, Etopophos®, VP-16)

Etoposide works by blocking the action of an enzyme in cells called topoisomerase, which is necessary for cell replication and tumor growth. Cells need this enzyme to keep their DNA in the proper shape when they are dividing. Blocking this enzyme leads to breaks in the DNA, which leads to cancer cell death.

### How to Take Etoposide

Etoposide is given by intravenous (IV, into a vein) infusion. It can also be given by mouth in the form of a [capsule](#), but this article will focus on the intravenous route. The dosage and schedule is determined by the person's size, type of cancer, and mode of administration. It can be given alone or with other medications.

Even when carefully and correctly administered by trained personnel, this drug may cause a feeling of burning and pain. There is a risk that this medication may leak out of the vein at the injection site, resulting in tissue damage that can be severe. If the area of injection becomes red, swollen, or painful at any time during or after the injection, notify your care team immediately. Do not apply anything to the site unless instructed by your care team.

### Possible Side Effects of Etoposide

There are a number of things you can do to manage the side effects of etoposide. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

#### Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your [WBC count can drop](#), putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4°F or 38°C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

## Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bathe daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with has any vaccinations.

## Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing, or pain in your chest. If the count gets too low, you may receive a blood transfusion.

## Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib) etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding use of these agents and all over the counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

## Mouth Ulcers (Mucositis)

Certain cancer treatments can cause [sores or soreness in your mouth](#) and/or throat. Notify your oncology care team if your mouth, tongue, inside of your cheek, or throat becomes white, ulcerated, or painful. Performing [regular mouth care](#) can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt with warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon of salt in an eight ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages and citrus juices.

## Nausea and/or Vomiting

Talk to your oncology care team so they can prescribe medications to help you manage [nausea and vomiting](#).

In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

## Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become [thin, brittle, or may fall out](#). This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats, and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

## Fatigue

[Fatigue](#) is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

## Important but Less Common Side Effects

- **Lowering of Blood Pressure:** Blood pressure may drop while this medication is being infused. When receiving this medication through a vein, your nurse will be checking your blood pressure before and during the infusion. It may become necessary to stop the administration of this medication or slow down the infusion rate if your blood pressure drops.
- **Allergic Reactions:** Patients can have an allergic reaction to this medication. Signs of a reaction can include: swelling, chills, fever, increased heart rate, shortness of breath or difficulty breathing, or a decrease in blood pressure. If you notice any changes in how you feel during the infusion, let your nurse know immediately. The infusion will be slowed or stopped if this occurs.
- **Radiation Recall:** Radiation recall is when the administration of a medication causes a skin reaction that looks like a sunburn (redness, swelling, soreness, peeling skin) in areas where radiation was previously given. Notify your oncology team if you notice this side effect. Treatment can include topical steroid ointments and a delay in your next chemotherapy dose.
- **Secondary Malignancies:** There is a very low risk of developing leukemia due to treatment with this medication, which can occur many years after treatment. This is most often associated with repeated treatments or high doses.

## Sexual & Reproductive Concerns

This medication may affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and [vaginal dryness](#). In addition, the desire for sex may decrease during treatment.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment and for at least 6 months for women and 4 months for men after treatment, even if your menstrual cycle stops or you believe you are not producing sperm. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team. You should not breastfeed while receiving this medication.

## Prednisone (Sterapred®, Prednisone Intensol)

**Pronounce:** PRED-ni-sone

**Classification:** Glucocorticoid

## About Prednisone (Sterapred®, Prednisone Intensol)

Prednisone is a corticosteroid/glucocorticoid that is similar to a hormone made naturally in your body. Corticosteroids (sometimes called "steroids") are used to lessen inflammation (swelling and/or redness). Steroids may be used for a few diseases, such as asthma, autoimmune disorders, reactions to medications, and gastrointestinal (GI) disorders (colitis). Prednisone may be given to prevent a reaction to a medication, to prevent or decrease nausea, or it can be used in high doses to treat some types of cancers.

## How to Take Prednisone

Prednisone comes as a tablet to take by mouth. The tablet should be swallowed whole. You should not break or chew the tablet. Prednisone is best taken with food or milk, as it can upset your stomach. Unless told otherwise, take your dose(s) of prednisone at the same time(s) each day. Your dosing schedule depends on what the medication is being used for. If you miss a dose, take it as soon as possible. If it is too close to your next dose, skip the missed dose and go back to your normal schedule.

Do not stop taking this medication without first speaking to your care provider as this could cause side effects such as weakness, fatigue, decreased appetite, weight loss, nausea, vomiting, diarrhea, and abdominal (belly) pain.

This medication can affect the levels of many other medications including warfarin, cyclosporine, oral birth control pills, phenytoin, bupropion, thalidomide, erythromycin, ketoconazole, and ritonavir, among others. Be sure to tell your healthcare provider about all medications and supplements you take.

You, or anyone you live with, should avoid having live or live-attenuated vaccines while receiving this medication. These include herpes zoster (Zostavax) for shingles prevention, oral polio, measles, nasal flu vaccine (FluMist®), rotavirus, and yellow fever vaccines.

## Storage and Handling

Store your medication in the original, labeled container at room temperature and in a dry location (unless otherwise directed by your healthcare provider or pharmacist). Keep containers out of reach of children and pets.

## Where do I get this medication?

Prednisone is available through most pharmacies. Your oncology team will work with your prescription drug plan to identify an in-network pharmacy for the distribution of this medication.

## Insurance Information

This medication may be covered under your prescription drug plan. Patient assistance may be available to qualifying individuals without prescription drug coverage. Your care team can help you find these resources if they are available.

## Possible Side Effects of Prednisone

There are a number of things you can do to manage the side effects of prednisone. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

### Swelling

Patients may notice swelling in their hands and/or feet. Elevating (raising) the feet may help to lessen swelling in the feet and ankles. Avoid restrictive or tight clothing that may make it harder for the fluid to drain from the hands, feet, and ankles.

## Increased Risk of Infection

This medication can lower your ability to fight new and current infections. Contact your provider if you have any signs of infection, such as fever, chills, sore throat or cold, cough or burning with urination. Tell your provider if you have any recent or ongoing infections, including viral or bacterial. Tell your provider if you have had any recent vaccinations. Avoid being around anyone who has chickenpox or measles.

Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever, or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bathe daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your doctor or nurse before scheduling dental appointments or procedures.
- Ask your doctor or nurse before you, or someone you live with has any vaccinations.

## Weakening of the Bones (Osteoporosis)

Long-term use can lead to early osteoporosis. Your provider may have you get a [bone density scan \(DEXA scan\)](#) to check your bone health if you are on long-term therapy.

## Eye Problems

Long-term use of this medication can increase the pressure in your eyes. Report any changes in vision, blurry or double vision, and eye pain or redness.

## Increase in Appetite

Prednisone can cause you to be hungrier or thirstier than usual. Drink plenty of fluids and try to make your snacks healthy ones.

## Increase in Energy

Prednisone can cause an increase in energy. You may also have insomnia, or difficulty sleeping. Taking the medication in the morning may help prevent this.

## Irritability or Change in Mood

Some patients report feeling irritable or might notice a change in their mood while taking prednisone. If this becomes difficult to handle or if you feel like you may harm yourself or others, call your healthcare provider right away.

## Nausea and Heartburn

Taking prednisone with food or milk is usually enough to prevent nausea and heartburn. If possible, take the medication when you can be upright (not lying down) for a few hours after the dose. Avoid things that worsen the symptoms and try antacids (milk of magnesia and calcium tablets, like Tums), saltines, or ginger ale to lessen symptoms.

## Increased Blood Sugar

Prednisone can raise your blood sugar. Diabetics should watch their blood sugar closely and may need higher doses of insulin while taking prednisone. Patients who are not diabetic but have increased blood sugar levels may be told to check their blood sugar and take insulin while on prednisone. Your provider will tell you if this is necessary.

## GI Bleed & Tear

This medication can cause bleeding or a tear in your intestinal wall (GI tract). Signs of these problems may be: unexpected bleeding, blood in the stool or black stools, coughing up blood, vomiting blood, vomit that looks like coffee grounds, fever, severe pain in the abdomen, or new abdominal swelling. If you have any of these, contact your oncology care team right away or go to the emergency room.

## Other Side Effects

Prednisone can cause delayed wound healing, headaches, muscle weakness, and cataracts (after long-term use).

## Reproductive Concerns

Exposure of an unborn child to this medication could cause birth defects in rare cases, so you should not become pregnant or father a child while on this medication. Even if your menstrual cycle stops or you believe you are not producing sperm, you could still be fertile and conceive. You should check with your healthcare team before breastfeeding while receiving this medication.

# Procarbazine (Matulane®)

**Pronounce:** proe-KAR-ba-zeen

**Classification:** Alkylating Agent

## About Procarbazine (Matulane®)

Procarbazine exerts its anti-cancer effect through a process called alkylation. Alkylation damages the DNA of cells, which prevents them from dividing, and causes them to die. Since cancer cells, in general, divide faster and with less error-correcting than healthy cells, cancer cells are more sensitive to this damage.

## How to Take Procarbazine

Procarbazine comes as a capsule, to be taken by mouth. Take the capsule whole, do not break, chew, or open the capsules. The dosage and schedule are determined by the person's size. If you forget to take a dose, take it as soon as you remember. If it is close to your next dose do not double your dose, rather skip the dose and resume your usual dosing schedule.

It is important to make sure you are taking the correct amount of medication every time. Before every dose, check that what you are taking matches what you have been prescribed.

Patients should avoid alcoholic beverages while taking procarbazine. The combination of alcohol and procarbazine can cause unpleasant side effects. Foods high in tyramine (wine, ripe cheeses, and bananas) should also be avoided. Consuming foods high in tyramine while taking procarbazine can raise your blood pressure. Numerous medications including barbituates, antihistamines, narcotics, and phenothiazines can cause central nervous system depression and should be avoided. Make sure your provider is aware of all the medications, vitamins, and supplements you are taking. Patients should avoid smoking while taking procarbazine because there is an increased risk of developing lung cancer.

## Storage and Handling



Store your medication in the original, labeled container at room temperature and in a dry location (unless otherwise directed by your healthcare provider or pharmacist). This medication should not be stored in a pillbox. Keep containers out of reach of children and pets.

If a caregiver prepares your dose for you, they should consider wearing gloves or pour the pills directly from their container into the cap, a small cup, or directly into your hand. They should avoid touching the pills. They should always wash their hands before and after giving you the medication. Pregnant or nursing women should not prepare the dose for you. Ask your oncology team where to return any unused medication for disposal. Do not flush down the toilet or throw it in the trash.

## Where do I get this medication?

Procarbazine is available through select specialty pharmacies. Your oncology team will work with your prescription drug plan to identify an in-network specialty pharmacy for the distribution of this medication and shipment directly to your home.

## Insurance Information

This medication may be covered under your prescription drug plan. Patient assistance may be available to qualifying individuals without prescription drug coverage. Co-pay cards, which reduce the patient co-pay responsibility for eligible commercially (non-government sponsored) insured patients, are also available. Your care team can help you find these resources if they are available.

## Possible Side Effects of Procarbazine

There are a number of things you can do to manage the side effects of procarbazine. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common side effects:

### Nausea and/or Vomiting

Talk to your oncology care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy, or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

### Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your [WBC count can drop](#), putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4°F or 38°C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

### Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever, or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bathe daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.

- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with has any vaccinations.

## Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing, or pain in your chest. If the count gets too low, you may receive a blood transfusion.

## Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums, or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib), etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding the use of these agents and all over-the-counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

## Mouth Ulcers (Mucositis)

Certain cancer treatments can cause [sores or soreness in your mouth](#) and/or throat. Notify your oncology care team if your mouth, tongue, inside of your cheek, or throat becomes white, ulcerated, or painful. Performing [regular mouth care](#) can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt with warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon of salt in an eight-ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages, and citrus juices.

## Less common, but important side effects

- **Secondary Malignancies:** There is a slight risk of developing leukemia or another type of cancer due to treatment with this medication. This is most often associated with repeated treatments or high doses. Patients should avoid smoking if they have received procarbazine. There is an increased risk of developing lung cancer after treatment with this medication and smoking can further increase the risk.
- **Central Neurotoxicity:** this can present as weakness, loss of balance or coordination, unsteadiness, dizziness, or drowsiness. Notify your healthcare provider right away if you experience any of these symptoms.
- **Allergic reaction:** In some cases, patients can have an allergic reaction to this medication. Signs of a reaction can include shortness of breath or difficulty breathing, chest pain, rash, flushing or itching, or a decrease in blood pressure. Inform your care team immediately if you have symptoms of an allergic reaction.
- **Diarrhea:** Your oncology care team can recommend medications to relieve [diarrhea](#). Also, try eating

[low-fiber](#), bland foods, such as white rice and boiled or baked chicken. Avoid raw fruits, vegetables, whole-grain breads, cereals, and seeds. Soluble fiber is found in some foods and absorbs fluid, which can help relieve diarrhea. Foods high in soluble fiber include applesauce, bananas (ripe), canned fruit, orange sections, boiled potatoes, white rice, products made with white flour, oatmeal, cream of rice, cream of wheat, and farina. Drink 8-10 glasses of non-alcoholic, uncaffeinated fluid a day to prevent dehydration.

- **Peripheral Neuropathy (Numbness or Tingling in the Hands and/or Feet):** [Peripheral neuropathy](#) is a toxicity that affects the nerves. It causes numbness or a tingling feeling in the hands and/or feet, often in the pattern of a stocking or glove. This can get progressively worse with additional doses of the medication. In some people, the symptoms slowly resolve after the medication is stopped, but for some, it never goes away completely. You should let the oncology care team know if you experience numbness or tingling in the hands and/or feet, as they may need to adjust the doses of your medication.

## Sexual and Reproductive Changes

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment, even if your menstrual cycle stops or you believe you are not producing sperm. You should consult with your healthcare team before breastfeeding while receiving this medication.

This medication may affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and [vaginal dryness](#). In addition, the desire for sex may decrease during treatment. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team.

## Vincristine (Oncovin®, Vincasar PFS®, VCR)

**Pronounce:** vin-KRIS-teen

**Classification:** Antimicrotubule Agent

### About Vincristine (Oncovin®, Vincasar PFS®, VCR)

Vincristine is a member of the vinca alkaloids family of chemotherapy agents. These medications work by interfering with cell division, which leaves the tumor unable to grow and spread. Vincristine was developed from the periwinkle plant.

### How to Take Vincristine

This medication is administered intravenously (IV, into a vein) by a trained professional. Your dose and treatment schedule depend on your size and the type of cancer being treated.

This medication is a vesicant. Even when carefully and correctly administered by trained personnel, this drug may cause a feeling of burning and pain. There is a risk that this medication may leak out of the vein at the injection site, resulting in tissue damage that can be severe. If the area of injection becomes red, swollen, or painful at any time during or after the injection, notify your doctor or nurse immediately. Do not apply anything to the site unless instructed by your doctor or nurse.

The blood levels of this medication can be affected by certain foods and medications, so they should be avoided. These include grapefruit, grapefruit juice, ketoconazole, rifampin, phenytoin, St. John's wort, and many anti-fungal medications. Be sure to tell your healthcare provider about all medications and supplements you take.

## Possible Side Effects of Vincristine

There are a number of things you can do to manage the side effects of vincristine. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

### Constipation

Vincristine can cause serious constipation, abdominal pain and can even lead to a blockage or stoppage of the bowel (called paralytic ileus) if not treated promptly. There are several things you can do to prevent or relieve constipation. Include fiber in your diet (fruits and vegetables), drink 8-10 glasses of non-alcoholic fluids a day, and keep active. A stool softener once or twice a day may prevent constipation. If you do not have a bowel movement for 2-3 days, you should contact your healthcare team for [suggestions to relieve the constipation](#).

### Neurotoxicity

This is a toxicity that affects the nerves. The most common effect is called peripheral neuropathy, which affects the nerves in the hands and feet, causing causes numbness or tingling, often in the pattern of a stocking or glove. This can get worse with additional doses of the medication and can lead to difficulty with balance or walking. In some people, the symptoms slowly get better after the medication is stopped, but for some, it never goes away completely. Let your healthcare provider know if you have numbness or tingling in the hands and feet, as they may need to adjust the doses of your medication.

The vinca alkaloid class of chemotherapies is known to cause neuropathy, but vincristine can also cause neurologic toxicity that presents as mental depression, headache, malaise, dizziness, and seizures. It can also cause toxicity of the cranial nerves, which affects the vocal cords (changes in voice), eyes (visual changes), or facial nerves (drooping of the face or mouth). Patients can develop severe pain in the jaw soon after the first treatment with vincristine, which is caused by the medication affecting the nerves. If you notice any of these problems, notify your healthcare team right away.

### Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your [WBC count can drop](#), putting you at a higher risk of getting an infection. You should let your provider know right away if you have a fever (temperature greater than 100.4°F or 38°C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

#### Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever, or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bathe daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with has any vaccinations.

### Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing, or pain in your chest. If the count gets too low, you may receive a blood

transfusion.

## Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums, or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib), etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding the use of these agents and all over-the-counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

## Mouth Ulcers (Sores)

Certain cancer treatments can cause [sores or soreness in your mouth](#) and/or throat. Notify your oncology care team if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated, or painful. Performing [regular mouth care](#) can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt with warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon of salt in an eight-ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages, and citrus juices.

## Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become [thin, brittle, or may fall out](#). This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats, and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

## Fatigue

[Fatigue](#) is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

## Nausea and/or Vomiting

Talk to your care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try antacids, (e.g. milk of magnesia, calcium tablets such as Tums), saltines, or ginger ale to lessen symptoms.

Call your doctor or nurse if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

## Important but Less Common Side Effects

- **Urinary retention:** This is the inability to urinate even when you feel that you need to. If you notice you are unable to urinate, you should call your healthcare team right away or go to the emergency room to be evaluated.
- **Allergic reaction:** Some patients will have an allergic reaction to the medication during the infusion or shortly after. Let your nurse know right away if you have any shortness of breath or difficulty breathing, rash, or swelling of the face.
- **Radiation Recall:** This medication may cause radiation recall. It may present as a skin reaction that looks like a sunburn (redness, swelling, soreness, peeling skin) in areas where radiation was previously given. Notify your oncology care team if you notice this side effect. Treatment can include topical steroid ointments and a delay in your next chemotherapy dose.

## Sexual & Reproductive Concerns

This drug may affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and [vaginal dryness](#). In addition, the desire for sex may decrease during treatment.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment, even if your menstrual cycle stops or you believe you are not producing sperm. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team. You should consult with your healthcare team before breastfeeding while receiving this medication.

---

OncoLink is designed for educational purposes only and is not engaged in rendering medical advice or professional services. The information provided through OncoLink should not be used for diagnosing or treating a health problem or a disease. It is not a substitute for professional care. If you have or suspect you may have a health problem or have questions or concerns about the medication that you have been prescribed, you should consult your health care provider.