



## Cladribine (2-CDA, Leustatin®)

**Pronounce:** KLAD-rih-bean

**Classification:** Antimetabolite

### About Cladribine (2-CDA, Leustatin®)

Cladribine is a type of chemotherapy called an antimetabolite. These medications help kill cancer cells by affecting the cell's DNA.

### How to Take Cladribine

Cladribine is given by intravenous (IV, into a vein) infusion. The dose depends on your body size and type of cancer. If you develop redness, swelling, or soreness at the injection site, tell a member of your care team right away.

### Possible Side Effects of Cladribine

There are a number of things you can do to manage the side effects of cladribine. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

#### Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your [WBC count can drop](#), putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4°F or 38°C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

#### Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever, or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with has any vaccinations.

#### Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness

of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion.

## Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums, or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib), etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding the use of these agents and all over-the-counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

## Neurotoxicity

There have been reports of this medication affecting the nervous system. Notify your provider if you develop weakness, paralysis, confusion, you are difficult to arouse or develop pins and needles in your fingers/toes.

## Kidney Problems

This medication can cause kidney problems, including an increased creatinine level, which your oncology care team may monitor for using blood tests. Notify your healthcare provider if you notice decreased urine output, blood in the urine, swelling in the ankles, or loss of appetite.

## Fever

Fever can occur after the dose of cladribine is given. Fever is most often seen when treating hairy cell leukemia. You should always call your care team if you develop a fever greater than 100.4°F or 38°C, as it could be caused by an infection. You can have a fever even if your white blood cell count is not low.

## Fatigue

[Fatigue](#) is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

## Nausea and/or Vomiting

Talk to your oncology care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

## Important but Less Common Side Effects

- **[Tumor Lysis Syndrome](#):** If there are many tumor cells in your body prior to treatment, you are at risk for tumor lysis syndrome. This happens when the tumor cells die too quickly and their waste overwhelms the body. You may be given a medication (allopurinol) and IV fluids to help prevent this. If you have nausea, vomiting, diarrhea, or become lethargic (drowsy, sluggish), call your oncology team

right away. TLS can affect your kidney function. Your provider will monitor your kidney function with blood work. Tell your provider if you have little or no urine output.

## Live Vaccines

You, or anyone you live with, should avoid having live or live-attenuated vaccines while receiving this medication. These include herpes zoster (Zostavax) for shingles prevention, oral polio, measles, nasal flu vaccine (FluMist®), rotavirus, and yellow fever vaccines.

## Reproductive Concerns

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment. Even if your menstrual cycle stops or you believe you are not producing sperm, you could still be fertile and conceive. You should not breastfeed while receiving this medication.

---

OncoLink is designed for educational purposes only and is not engaged in rendering medical advice or professional services. The information provided through OncoLink should not be used for diagnosing or treating a health problem or a disease. It is not a substitute for professional care. If you have or suspect you may have a health problem or have questions or concerns about the medication that you have been prescribed, you should consult your health care provider.