



## Mechlorethamine (Mustargen®, Nitrogen Mustard)

**Pronounce:** me-klor-ETH-a-meen

**Classification:** Alkylating Agent

### About Mechlorethamine (Mustargen®, Nitrogen Mustard)

\*\*\*This medication is no longer being used but you may have received it as part of your cancer treatment.\*\*\*

Mechlorethamine is a member of a class of chemotherapies called alkylating agents. It used to be called nitrogen mustard, and you may still hear it called this. It is a derivative of nitrogen gas and was first used in World War I as a chemical warfare agent. It was found to have an effect on bone marrow and white blood cells, so research began to investigate its use as a treatment for lymphoma.

Mechlorethamine exerts its anti-cancer effect by a process called alkylation. Alkylation damages the DNA of cells, which prevents them from dividing, and causes them to die. Since cancer cells, in general, divide faster and with less error-correcting than healthy cells, cancer cells are more sensitive to this damage. In this way, mechlorethamine slows or stops the growth of cancer cells in your body.

### How to Take Mechlorethamine

Mechlorethamine is given by intravenous (IV, into a vein) injection. The dosage and schedule are determined by the person's size and type of cancer. In addition, mechlorethamine can be delivered directly into body cavities (lining of the lung, abdomen, and heart).

Even when carefully and correctly administered by trained personnel, this drug may cause a feeling of burning and pain. There is a risk that this medication may leak out of the vein at the injection site, resulting in tissue damage that can be severe. If the area of injection becomes red, swollen, or painful at any time during or after the injection, notify your care team immediately. Do not apply anything to the site unless instructed by your care team.

### Possible Side Effects of Mechlorethamine

There are a number of things you can do to manage the side effects of mechlorethamine. Talk to your care team about these recommendations. They can help you decide what will work best for you. These are some of the most common or important side effects:

#### Nausea and/or Vomiting

Talk to your oncology care team so they can prescribe medications to help you manage [nausea and vomiting](#). In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy, or acidic foods (lemons, tomatoes, oranges). Try saltines, or ginger ale to lessen symptoms.

Call your oncology care team if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

#### Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your [WBC count can](#)

[drop](#), putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4°F/38°C), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

### Tips to preventing infection:

- [Washing hands](#), both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever, or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bathe daily and perform frequent [mouth care](#).
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your oncology care team before scheduling dental appointments or procedures.
- Ask your oncology care team before you, or someone you live with has any vaccinations.

### Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the [red cell count is low](#), you may feel tired or weak. You should let your oncology care team know if you experience any shortness of breath, difficulty breathing, or pain in your chest. If the count gets too low, you may receive a blood transfusion.

### Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the [count is low](#) you are at a higher risk of bleeding. Let your oncology care team know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums, or blood in your urine or stool. If the platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin/Advil (ibuprofen), Aleve (naproxen), Celebrex (celecoxib), etc. as these can all increase the risk of bleeding. Please consult with your healthcare team regarding the use of these agents and all over-the-counter medications/supplements while on therapy.
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

### Fatigue

[Fatigue](#) is very common during cancer treatment and is an overwhelming feeling of exhaustion that is not usually relieved by rest. While on cancer treatment, and for a period after, you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team for helpful tips on dealing with this side effect.

### Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become [thin, brittle, or may fall out](#). This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats, and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun.

## Decrease in Appetite

**Nutrition** is an important part of your care. Cancer treatment can affect your appetite and, in some cases, the side effects of treatment can make eating difficult. Ask your oncology care team about nutritional counseling services at your treatment center to help with food choices.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.
- You may experience a metallic taste or find that food has no taste at all. You may dislike foods or beverages that you liked before receiving cancer treatment. These symptoms can last for several months or longer after treatment ends.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products, and fish without a strong smell. Sometimes cold food has less of an odor.
- Add extra flavor to meat or fish by marinating it in sweet juices, sweet and sour sauce, or dressings. Use seasonings like basil, oregano, or rosemary to add flavor. Bacon, ham, and onion can add flavor to vegetables.

## Secondary Cancers

A secondary cancer is one that develops as a result of cancer treatment for another cancer. This is quite rare, but you should be aware of the risk. In most cases, a secondary cancer related to chemotherapy is a blood cancer (leukemia, lymphoma). This can occur years after treatment. This is most often associated with repeated treatments or high doses. Your provider will monitor your labs closely. Consider having a complete blood count with differential checked annually by your healthcare provider if you received high-risk therapies.

## Hypersensitivity and Allergic Reaction

In some cases, patients can have hypersensitivity or an allergic reaction to this medication. Signs of a reaction can include shortness of breath or difficulty breathing, chest pain, rash, flushing or itching, or a decrease in blood pressure. If you notice any changes in how you feel during the infusion, let your nurse know immediately.

## Skin Changes

Some patients may develop a rash, very dry skin, or blisters. You may notice a darkening of the veins used for infusion. Use an alcohol-free moisturizer on your skin and lips; avoid moisturizers with perfumes or scents. Your doctor or nurse can recommend a topical medication if itching is bothersome. If your skin does crack or bleed, be sure to keep the area clean to avoid infection. Be sure to notify your healthcare provider of any rash that develops, as this can be a reaction. They can give you more tips on [caring for your skin](#).

## Less common but important side effects can include:

- **Tumor Lysis Syndrome:** If there are a large number of tumor cells in your body prior to treatment, you are at risk for tumor lysis syndrome. This happens when the tumor cells die too quickly and their waste overwhelms the body. You may be given a medication (allopurinol) and IV fluids to help prevent this. If you experience nausea, vomiting, diarrhea or become lethargic (drowsy, sluggish), notify your oncology team right away. TLS can affect your kidney function. Your provider will monitor your kidney function with blood work. Notify your provider if you have little or no urine output.
- **Vertigo:** Vertigo is ringing in the ears or diminished hearing. Report symptoms of dizziness or hearing changes to your healthcare team.

## Sexual & Reproductive Concerns

This medication may affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes

and [vaginal dryness](#). In addition, the desire for sex may decrease during treatment. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment even if your menstrual cycle stops or you believe you are not producing sperm. You should consult with your healthcare team before breastfeeding while receiving this medication.

---

OncoLink is designed for educational purposes only and is not engaged in rendering medical advice or professional services. The information provided through OncoLink should not be used for diagnosing or treating a health problem or a disease. It is not a substitute for professional care. If you have or suspect you may have a health problem or have questions or concerns about the medication that you have been prescribed, you should consult your health care provider.